



Letter to the Editor

Incidentally Detected Meckel Diverticulum during Cystectomy: An Unhinging Dilemma?

Manjeet Kumar¹

¹Department of Urology, Indira Gandhi Medical College, Shimla, Himachal Pradesh, India.

***Corresponding author:**

Manjeet Kumar, MBBS, MS,
MCh, Department of Urology,
IGMC, Shimla, HP, India.

manjeetkumar.1014@gmail.com

Epub Ahead of Print:

4 April 2023

Published: 30 October 2024

DOI

10.1055/s-0043-1761468

Dear Editor,

Meckel's diverticulum is found in 2% of the population, which makes it the commonest congenital abnormality of the gastrointestinal tract. It is diagnosed as an incidental finding during surgery for other reasons. It is found 100 cm from the ileocecal junction, at the antimesenteric border. Other presentations can be abdominal pain, infection, bleeding from ectopic mucosa, recurrent intestinal obstruction, etc. It is formed during the embryonic period due to incomplete obliteration of the omphalomesenteric duct. Treatment of this complication is the removal of Meckel's diverticulum.^[1,2]

Incidentally found Meckel's diverticulum during urological surgeries is a surgical dilemma. Complications due to the removal of Meckel's diverticulum could be challenging to manage, at least as mentioned in earlier studies.^[3]

Cullen et al in a study reported 6.4% chances of developing complications due to Meckel's diverticulum on long-term follow-up. When diverticulectomy was done, mortality and morbidity of 2% and 12%, respectively, were noted in these patients. In contrast, mortality and morbidity in incidental diverticulectomy were 1% and 2% respectively. On further analysis, it was found that complications in the incidental surgery group were related to the primary operation or patient fitness rather than diverticulectomy itself.^[4]

Park et al recommended resection of Meckel's in male patients, diverticulum length more than 2 cm, abnormal thickening, or appearance of a diverticulum.^[1] Carcinoid tumors were found in Meckel's diverticulum in 2.1% of asymptomatic patients and 2.2% for symptomatic ones.^[5,6]

In the literature, it is clear, especially in the recent literature, there is a move toward resection for all or in those with high-risk factors.^[4]

Regarding incidental detection of Meckel's during urological surgeries, a few case reports are available. Meckel's has been used during ureteric repair to bridge gap as a part of Mitrofanoff procedure.^[7] In other case report Meckel's has been used to implant ureter after radical cystectomy to the Bricker segment.^[8] Another case report used Meckel's as part of orthotopic neobladder to lengthen the distal part for urethral anastomosis.^[9]

Although there is no compelling evidence in the current literature to support prophylactic diverticulectomy for patients with MD, we believe that the removal of Meckel's should be done

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

©2024 Published by Scientific Scholar on behalf of International Journal of Recent Surgical and Medical Sciences



Figure 1: Incidentally detected Meckel's during radical cystectomy.

in young patients, with suspicious thickness of more than 2 cm diverticulum, especially in patients undergoing radical cystectomy.

In our experience, we encountered two incidental Meckel's diverticulum during 86 radical cystectomies [Figure 1]. In one patient, Meckel's was removed as it was located more distally near the ileocecal junction. In another patient, it was left in situ as no abnormality was found. No complications were seen in both patients related to Meckel's diverticulum. In our view, Meckel's should be removed if Meckel's is abnormal, thickened, more than 2 cm, and near the intestinal anastomosis. Complications related to Meckel's are more difficult to diagnose and manage after radical cystectomy, if they occur, unfortunately.

Funding

None.

Conflicts of interest

None declared.

REFERENCES

1. Park JJ, Wolff BG, Tollefson MK, Walsh EE, Larson DR. Meckel Diverticulum: The Mayo Clinic Experience with 1476 Patients (1950-2002). *Ann Surg* 2005;241:529-33.
2. Sagar J, Kumar V, Shah DK. Meckel's Diverticulum: A Systematic Review. *J R Soc Med* 2006;99:501-5.
3. Leijonmarck CE, Bonman-Sandelin K, Frisell J, Räf L. Meckel's Diverticulum in the Adult. *Br J Surg* 1986;73:146-9.
4. Cullen JJ, Kelly KA, Moir CR, Hodge DO, Zinsmeister AR, Melton L III. Surgical Management of Meckel's Diverticulum. An Epidemiologic, Population-based Study. *Ann Surg* 1994;220:564-8, discussion 568-9
5. Silk YN, Douglass HO Jr, Penetrante R. Carcinoid Tumor in Meckel's Diverticulum. *Am Surg* 1988;54:664-7.
6. Bona D, Schipani LS, Nencioni M, Rubino B, Bonavina L. Laparoscopic Resection for Incidentally Detected Meckel Diverticulum. *World J Gastroenterol* 2008;14:4961-3.
7. Adams J, Djakovic N, Gilfrich C, *et al.* Ureteric Replacement with Meckel's Diverticulum. *BJU Int* 2007;99:647-50.
8. Wyczółkowski M, Klima W, Kotuła J, Malik L. Meckel's Diverticulum as the Site of Implanting Ureter to the Bricker Segment After Radical Cystectomy. *J Urol* 2003;169:285.
9. Van Haute W, Ghysel C, Van Oyen P. The Use of a Meckel's Diverticulum in the Creation of an Orthotopic Neobladder in Case of a Short Mesoileum: A Case Report. *Adv Urol* 2009;2009:493236.

How to cite this article: Kumar M. Incidentally Detected Meckel Diverticulum during Cystectomy: An Unhinging Dilemma? *Int J Recent Sur Med Sci.* 2024;10:S111-2. doi: 10.1055/s-0043-1761468